




Five Years Experience of Peripheral Veno-Arterial ECMO as Method of Mechanical Circulatory Support in Heart Transplant Candidates

V. N. Poptsov, E. A. Spirina, A. A. Dogonascheva.

*Heart/Lung Transplant Team,
Shumakov National Medical Research Centre of Transplantology and Artificial Organs,
Moscow, Russian Federation*

Adult Heart Transplants % of Patients Bridged with Mechanical Circulatory Support* by Year and Device Type






Short-term mechanical circulatory support as a bridge to urgency heart transplantation

Spanish National Heart Transplant Registry

(Spanish, 2000-2009 yrs.)


Short-term pretransplant MCS



Barge-Caballero E. et al. Impact of short-term mechanical circulatory support with extracorporeal devices on postoperative outcomes after emergency heart transplantation: **data from a multi-institutional Spanish cohort.** // *Int J Cardiol* 2014; 173: 86-93.

High-Urgency Waiting List Super Urgence (SU) (France, since 2004 yr.)


38% (n=87) **SU1** with pretransplant VAECMO



D'Alessandro C. et al. High-urgency waiting list for cardiac recipients in France: **single-centre 8-year experience.** // *Eur J Cardiothorac Surg* 2017; 51: 271-278.

Shumakov National Medical Research Centre of Transplantology and Artificial Organs, Moscow

heart transplant wait list (2008-2017 yrs.)






Shumakov National Medical Research Centre of Transplantology and Artificial Organs, Moscow

heart transplant wait list

status UNOS 1A

(2008-2017 yrs.)





The goal of study was estimate results of using pVAECMO as a method of short-term mechanical circulatory support (MCS) in heart transplant candidates requiring urgent heart transplantation (HT).



Patients (n=149)

or **21.1%** (n=149) from heart transplant waiting list (n=706) (2011–2017 yrs.)


age 12-72 (43.0 ± 1.2) yrs

men 87% (n=129)/ female 13% (n=20)

TPG 4-20 (11.2 ± 2.0) mmHg


PVR 1.9-5.6 (3.8 ± 0.6) Wood's Unit

Etiology of advanced CHF



- dilated cardiomyopathy (n=95)
- coronary artery disease (n=42)
- restrictive cardiomyopathy (n=1)
- hypertrophic cardiomyopathy (n=1)
- peripartum cardiomyopathy (n=3)
- congenital or acquired heart disease (n=3)
- chronic cardiac allograft dysfunction (n=4)

INTERMACS



- level I
- level II

Patients (n=149)



**“conventional” VAECMO installation
(n=144 (96.6%))**

**extracorporeal cardiopulmonary
resuscitation (ECPR) at heart
transplant candidates with in-
hospital cardiac arrest
(n=5 (3.4%))**



peripheral VAECMO n=149 (100%)

open (surgical) technique
n=29 (19.5%)



percutaneous technique
n=120 (80.5%)




Leg's perfusion 100% (n=149) - required condition !!!

pVAECMO and left heart decompression (n=31 (20.8%))

percutaneous LA drainage (n=24 (72,7%))


open LV drainage (n=7 (21,2%))



1.72 ± 0.12 л/мин




3.83 ± 0.35 л/мин ($p < 0,05$)





bridge to transplantation (n=135 (90.6%))



VAECMO duration - from 12 h to 34 (5.8±3.2) days



parameter	M±σ
revolution per minute (rpm)	3719±102
Q, L/m	3.59±0.12
Q, L/min/m ²	1.84±0.10
sweep gas, L/min	3.2±0.4
sweep gas, FiO ₂	0.67±0.03



patients (n=149)

**conventional VAECMO installation
(n=144)**

**extracorporeal cardiopulmonary
resuscitation (ECPR) at heart transplant
candidates with in-hospital cardiac arrest
(n=5)**

**bridge to transplantation
(n=131)**

**death
(n=13)**

**death
(n=1)**

**bridge to transplantation
(n=4)**

death (n=14 (9.4%))

bridge to transplantation (n=135 (90.6%))



causes of death (n=14 (9.4%))

- ☐ multiorgan failure/sepsis (n=11 (78.6%))
- ☐ anoxic encephalopathy (n=1 (7.1%)) due to cardiopulmonary resuscitation
- ☐ stroke (n=2 (14.2%)) (preexisting LV thrombosis)



mortality risk factor analysis

variable	Odds Ratio (OR)	Confidence Interval	p Value
hemolysis ≥ 300 mg%	88.667	15.368-511.56	0.0040
pneumonia	33.156	6,94-156,92	0.0001
ALT ≥ 300 U/L	22.541	6.324-80.053	0.0001
AST ≥ 300 U/L	18.45	5.318-64.004	0.0001
total bilirubine ≥ 80 μ mol/l	21.607	5.977-78.110	0.0013
LV thrombosis	18.136	2.734-120.34	0.0060
transthoracic LV drainage	17.600	3.45-32.28	0.0015
BMI < 20 kg/m ²	13.1	2.842-60.389	0.0028
Creatinine ≥ 140 mmol/l	6.875	2.028-23.309	0.0013
urea ≥ 15 mmol/l	10.476	2.76-39.758	0.0002
cardiopulmonary resuscitation	2.519	0.2616-24.25	0.3938
percutaneous LV drainage	3.385	0.3276-34.939	0.3290
open (surgical) technique of cannulation	0.666	0.1407-3.159	1.0000



Heart transplantation at recipients with pretransplant pVAECMO (n=135)

hospital mortality **14,0%** (n=25)

donor risk factor

variable	Odds Ratio (OR)	Confidence Interval	p Value
donor age > 55 year	6,875	2,028-23,309	0,0013
noradrenalin > 600 ng/kg	10,476	2,76-39,758	0,0002
total bilirubine $\geq 80 \mu\text{mol/l}$	21,607	5,977-78,11	0,0010

Posttransplant Survival rate


Discharge to home	86.0%
1-year	84.5%
2-year	81.3%
3-year	78.1%

peritransplant risk factor

variable	Odds Ratio (OR)	Confidence Interval	p Value
VAECMO after HT > 4 day	21,4	2,392-191,46	0,0019
Inotropic index > 20	4,922	1,317-18,393	0,0234
bloodloss > 2.5 L	8,556	2,391-30,619	0,0234
packed red blood cell > 6	2,833	1,119-7,173	0,0378
fresh frozen plasma > 10	2,964	1,16-7,573	0,0244

Schumakov National Medical Research Center

310 (28.1%) HT at recipients with pretransplant VAECMO from 1102 HT (2011 – 2019 yrs)



Heart transplantation



Conclusion

Peripheral VA ECMO provides a successful bridge to urgent HT in 90% of heart transplant candidates with satisfactory survival after HT