

Early IVUS-guided selection for non-invasive cardiac allograft vasculopathy (CAV) screening in heart transplant recipients

L.M. Nelson MD¹, K. Rossing MD PhD DMSci¹, S. Boesgaard MD DMSci¹, F. Gustafsson MD PhD DMSci¹

¹Department of Cardiology, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark

ISHLT 2020 • Anniversary Meeting and Scientific Sessions



Relevant Financial Relationship Disclosure Statement

**Early IVUS-guided selection for non-invasive cardiac allograft vasculopathy (CAV)
screening in heart transplant recipients**


Lærke Marie Nelson, MD

I will not discuss off label use and/or investigational use of drugs/devices.


The following relevant financial relationships exist related to this presentation:

L.M. Nelson: No relationships to disclose. **K. Rossing:** Speaker fee from Abbott. **S. Boesgaard:** No relationships to disclose. **F. Gustafsson:** Scientific Medical Advisor for Novartis, Carmat, Abbot, and Pharmacosmos; Speaker fee from Novartis, Carmat, and Abbott.


Background – Survival in Heart Transplantation (HTx)



Background – Causes of Death in HTx




Background – Cardiac Allograft Vasculopathy (CAV)



- Vascular disease confined to cardiac allograft
- Intimal thickening → luminal narrowing
- Diffuse, circumferential, longitudinal
- Causing graft failure and death
- Early diagnosis crucial (prevention/treatment)
 - Yearly surveillance


Background – CAV Diagnosis & Surveillance

Coronary angiography




- ISHLT diagnostic criteria¹
- Risks (invasive, nephropathy)
- Underestimate CAV burden

Intravascular ultrasound (IVUS)



- Maximal intimal thickness (MIT)
- Subangiographic CAV (early)
- Not routinely used

Dobutamine stress echocardiography (DSE)



- Non-invasive modality
- High sensitivity CAV
- Low risk patients / renal function↓

¹Mehra MR et al. J Heart Lung Transplant 2010 Jul;29(7):717-27; Gustafsson F. UpToDate 2013.


Background – CAV Diagnosis & Surveillance

Coronary angiography




- ISHLT diagnostic criteria¹
- Risks (invasive, nephropathy)
- Underestimate CAV burden

Intravascular ultrasound (IVUS)



- Maximal intimal thickness (MIT)
- Subangiographic CAV (early)
- Not routinely used

Dobutamine stress echocardiography (DSE)



- Non-invasive modality
- High sensitivity CAV
- Low risk patients / renal function↓


Optimal strategy and risk stratification ???

Objective & Hypothesis

OBJECTIVE	Evaluate a strategy of early IVUS-guided selection to non-invasive screening for CAV in HTx recipients
HYPOTHESIS	Early IVUS-guided selection causes low risk of missing significant CAV as assessed by invasive coronary angiography at 5-year follow-up post-HTx


Methods - Algorithm

- Algorithm for early IVUS-guided selection to non-invasive CAV screening
- Employed at Rigshospitalet since 2009




Methods – Design & Study Population

- Single center, retrospective study
- Consecutive adult HTx recipients at Rigshospitalet from 2008-2018 following the screening algorithm
- N = 66



Results - Baseline

Table 1. Baseline characteristics (n=66)




■ low-risk group MIT < 0.5
■ high-risk group MIT ≥ 0.5

	All (n=66)	MIT < 0.5 (n=41)	MIT ≥ 0.5 (n=25)	p-value
Recipient characteristics				
Recipient age (years)	51 (33-60)	52 (33-59)	49 (42-63)	0.39
Male sex (n, %)	41 (62%)	25 (61%)	16 (64%)	0.81
LVAD (n, %)	26 (39%)	18 (44%)	8 (32%)	0.34
BMI (kg/m²)	25 (22-27)	24 (23-27)	25 (22-30)	0.26
Diabetes (%)	9 (9%)	5 (12%)	1 (4%)	0.27
HTx primary indication				
Idiopathic DCM (n, %)	36 (54%)	23 (56%)	13 (52%)	0.31
Ischemic (n, %)	15 (23%)	7 (17%)	8 (32%)	
Other* (n, %)	15 (23%)	11 (27%)	4 (16%)	
Donor characteristics				
Donor age (years)	41 (32-47)	34 (26-42)	45 (41-49)	0.0001
Donor male sex (n, %)	37 (56%)	22 (54%)	15 (60%)	0.62
Ischemic time (min)	240 (161-300)	275 (183-300)	222 (140-300)	0.50

Data are reported as median (IQR) or absolute number (percentage) as appropriate. Groups are compared with qui-square, Fishers exact or Mann-Whitney U test as appropriate. MIT=maximal intimal thickness, LVAD= left ventricular assist device, BMI=body mass index, HTx=heart transplantation, DCM=dilated cardiomyopathy. *HCM, RCM, ARVC, non-compaction, valvular, chemotherapy induced, and congenital.

Results – Outcome

Figure 1. Survival free from angiographic CAV according to MIT risk group



No at risk							
MIT < 0.5 mm:	41	40	37	35	33	30	0
MIT ≥ 0.5 mm:	25	25	19	16	12	5	0

18 patients met the combined endpoint:

- 4 patients died from any cause
 - 2 deaths (low-risk group)
 - 2 deaths (high-risk group)
- Angiographic CAV detected in 14 patients
 - 1 CAV (low-risk group)
 - 13 CAV (high-risk group)

Results – Outcome

Table 2. Cardiac allograft vasculopathy in the subgroup completing invasive coronary angiography at 5 years follow-up post-HTx (n=27)

	All (n=27)	MIT < 0.5 (n=15)	MIT ≥ 0.5 (n=12)	p-value
Any degree of CAV* (n, %)	11 (41%)	1 (7%)	10 (83%)	<.0001
ISHLT CAV1 (n, %)	9 (33%)	-	9 (75%)	
ISHLT CAV2 (n, %)	-	-	-	
ISHLT CAV3 (n, %)	2 (8%)	1 (7%)	1 (8%)	

Data are reported as absolute number (percentage). Groups are compared with qui-square or Fishers exact test as appropriate.
CAV=cardiac allograft vasculopathy, MIT=maximal intimal thickness, ISHLT= international society for heart and lung transplantation.
*According to the 2010 ISHLT criteria.


- Angiographic CAV was rare in low-risk patients
- No re-HTx
- No revascularization

Limitations

- Single center retrospective evaluation
- Small study population
- Incomplete adherence to algorithm
- No re-evaluation of coronary angiograms

Conclusions & Perspectives

- Early IVUS-guided selection to non-invasive CAV screening in selected patients appears to be safe
- Holds promise as a novel strategy for early risk stratification and CAV surveillance after HTx



- Reduce burden of invasive procedures in *low risk* patients
- Intensify preventive interventions in *high risk* patients

Acknowledgements

Co-authors

Department of Cardiology, Rigshospitalet, Copenhagen, Denmark

- Kasper Rossing MD, PhD, DMSci
- Søren Boesgaard MD, DMSci
- Finn Gustafsson MD, PhD, DMSci

Heart Transplant Program at Rigshospitalet

- Department of Cardiology
- Cath lab Section 2014 (IVUS, coronary angiography)
- Echo lab Section 2142 (DSE)

Financial support

- The Danish Heart Foundation
- The Research Foundation of Rigshospitalet
- The Research Foundation of The Heart Center Rigshospitalet



Rigshospitalet



DANISH HEART FOUNDATION

Thank you for your attention!

